

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau


INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/82, A01H 5/00	A1	(11) International Publication Number: WO 95/23230 (43) International Publication Date: 31 August 1995 (31.08.95)
(21) International Application Number: PCT/NO95/00042 (22) International Filing Date: 23 February 1995 (23.02.95) (30) Priority Data: 9403512.8 24 February 1994 (24.02.94) GB (71)(72) Applicants and Inventors: OLSEN, Odd-Arne [NO/NO]; Tarnveien 16, N-1430 Ås (NO). KALLA, Roger [AU/AU]; Mowle Place, Weetangera, ACT 2641 (AU). LINNESTAD, Casper [NO/NO]; Ovre Linnestad, N-1540 Vestby (NO). (74) Agent: A/S BERGEN PATENTKONTOR; Strandgt. 191, N- 5000 Bergen (NO).	(81) Designated States: AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW, MX, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD, SZ, UG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>	

(54) Title: PROMOTER FROM A LIPID TRANSFER PROTEIN GENE

(57) Abstract

An expression system for at least the aleurone cells of a developing caryopsis or for at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or developing grain or plant (e.g. in the root, leaves and stem) is described. The expression system comprises a gene promoter fused to a GOI (gene of interest). In a preferred embodiment the expression system comprises the GOI fused to a modified Ltp1 gene promoter.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

PROMOTER FROM A LIPID TRANSFER PROTEIN GENE

The present invention relates to a promoter and to a construct comprising the same.

- 5 In particular the present invention relates to the use of a promoter for the expression of a gene of interest (GOI) in a specific tissue or tissues of a plant.

More in particular the present invention relates to a modified promoter for a lipid transfer protein (Ltp) gene known as the Ltp1 gene. The present invention also
10 relates to the application of this modified Ltp1 gene promoter to express a GOI in a specific tissue or specific tissues of a plant. For example, expression can be in either the aleurone layer or the scutellar epithelial layer of a monocotyledon, especially a transgenic cereal caryopsis (or grain), more especially a developing transgenic cereal caryopsis (or grain). Particular examples include expression in the scutellar epithelial
15 tissue or vascular tissue of a transgenic rice plant, in particular in the vascular bundles and tip of emerging shoots and roots, leaf veins and vascular bundles of stems.

A diagrammatic illustration of a developing caryopsis (or grain) is presented in Figure
20 1, which is discussed in detail later. In short, a typical developing caryopsis (or grain) comprises an endosperm component and an embryo component. The endosperm, which is the site of deposition of different storage products such as starch and proteins, supports the growth of the emerging seedling during a short period of time after germination. The embryo gives rise to the vegetative plant. These
25 components and aspects are further discussed in Bosnes *et al.* 1992 and Olsen *et al.* 1992.

The embryo component can be divided into a scutellum and an embryo axis. The scutellum can be sub-divided into an epithelial layer, which is usually one cell thick,
30 and an inner body of parenchyma cells. Likewise, the embryo axis can be sub-divided into a root component and a shoot component.

The endosperm component of mature grains can be divided into a peripheral layer of living aleurone cells surrounding a central mass of non-living starchy endosperm cells. The aleurone layer in barley is three cells thick. During caryopsis germination, the cells of the aleurone layer produce amyolytic and proteolytic enzymes that degrade the storage compounds into metabolites that are taken up and are used by the growing embryo.

Two aspects of aleurone cell biology that have been intensively studied are the genetics of anthocyanin pigmentation of aleurone cells in maize (McClintock, 1987) and the hormonal regulation of gene transcription in the aleurone layer of germinating barley caryopsis (Fincher, 1989). Using transposon tagging, several structural and regulatory genes in the anthocyanin synthesis pathway have been isolated and characterized (Paz-Ares *et al.*, 1987; Dellaporta *et al.*, 1988). In barley, α -amylase and β -glucanase genes that are expressed both in the aleurone layer and embryos of mature germinating caryopsis have been identified (Karrer *et al.*, 1991; Slakeski and Fincher, 1992). In addition, two other cDNAs representing transcripts that are differentially expressed in the aleurone layers of developing barley grains have been isolated. These are CHI26 (Lea *et al.*, 1991) and pZE40 (Smith *et al.*, 1992).

None of these references discloses expression of those gene products in specific cell types of developing grains of transgenic cereal plants or in the scutellar epithelial tissue or vascular tissue of a germinating rice seedling or a developing rice grain or rice plant.

In the life of a developing caryopsis (or grain), the embryo component of a dried caryopsis will imbibe water. The presence of water triggers the production of the hormone gibberellic acid in the embryo. In barley and other grass caryopsis, the embryo releases the gibberellic acid which in turn causes expression of a number of genes in the aleurone layer of the endosperm resulting in the production of a number of enzymes such as α -amylases, proteases and β -glucanases. Similar enzymes are also produced by expression of genes in the epithelial layer.

These degradative enzymes digest certain components of the developing caryopsis (or grain) to form sugars and amino acids.

5 For example, the α -amylases digest the starch store in the starchy endosperm, whereas the proteases digest the storage proteins and the β -glucanases digest the cell walls. The resultant sugars and amino acids cross the epithelial layer and trigger growth of the shoot and root of the embryo axis - i.e. start the germination process.

10 In some cases it is desirable to transform seeds, grains, caryopsis and plants by introducing genes which, as a result of their expression, yield new or improved properties to the resulting transformed seeds, grains, caryopsis or plants. For example, it may be desirable to alter the expression levels of a natural structural gene which may be under- or over- expressed. It may even be desirable to reduce or eliminate a disease which harms or destroys the seed, grain, caryopsis or the plant.

15

It may even be desirable to make the seed, grain, caryopsis or the plant resistant to herbicides. It may even be desirable to prevent or to reduce the extent of pre-harvest sprouting.

20

It may even be desirable for the seed, grain, caryopsis, or plant to produce compounds useful for mammalian usage, such as human insulin.

Some techniques are known for addressing some of those aims.

25

For example, the bacterium *Agrobacterium tumefaciens* has been used to introduce desired genes into the chromosome of a plant. For example the gene coding for EPSP synthase, a key enzyme in the synthesis of aromatic acids in plants, has been isolated and introduced into petunia plants under the control of a CaMV promoter
30 (Shah *et al.*, [1986]). The transgenic plants expressed increased levels of EPSP synthase in their chloroplasts and were more tolerant to glyphosphate - which inhibits production of EPSP synthase.

Other examples may be found in R.W. Old & S.B. Primrose (1993). Another use of *Agrobacterium tumefaciens* is described in De Silva *et al.* (1992) wherein a recombinant DNA construct is described containing a plant plastid specific promoter that expresses a gene placed under its control in concert with the fatty acid or lipid biosynthesis in the plant cell.

PCT WO 90/01551 mentions the use of the aleurone cells of mature, germinating caryopsis to produce proteins from GOIs under the control of an α -amylase promoter. This promoter is active only in germinating caryopsis.

10

Non-specific lipid transfer proteins (nsLtps) have the ability to mediate *in vitro* transfer of radiolabelled phospholipids from liposomal donor membranes to mitochondrial acceptor membranes (Kader *et al.*, 1984; Watanabe and Yamada, 1986). Although their *in vivo* function remains unclear, nsLTPs from plants have recently received much attention due to their recurrent isolation as cDNA clones representing developmentally regulated transcripts expressed in several different tissues. A common feature is that, at some point in their development, they are highly expressed in tissues producing an extracellular layer rich in lipids.

20 In particular, transcripts corresponding to cDNAs encoding 10 kDa nsLTPs have been characterized in the tapetum cells of anthers as well as the epidermal layers of leaves and shoots in tobacco (Koltunow *et al.*, 1990; Fleming *et al.*, 1992), and barley aleurone layers (Mundy and Rogers, 1986; Jakobsen *et al.*, 1989).

25 In addition, a 10 kDa nsLTP has been discovered to be one of the proteins secreted from auxin-treated somatic carrot embryos into the tissue culture medium (Stern *et al.*, 1991).

30 Based on *in situ* hybridisation data demonstrating that the Ltp transcripts are localized in the protoderm cells of the somatic and zygotic carrot embryo, it was suggested that *in vivo* nsLTPs are involved in either cutin biosynthesis or in the biogenesis and degradation of storage lipids (Sossountzov *et al.*, 1991; Stern *et al.*, 1991).

A nsLTP in *Arabidopsis* has been localized to the cell walls lending further support to an extracellular function of this class of proteins (Thoma *et al.*, 1993).

5 Recently, using a standard *in vitro* Ltp assay, two 10 kDa nsLtps and one member of a novel class of 7 kDa nsLtp's were isolated from wheat seeds (Monnet, 1990; Dieryck *et al.*, 1992).

The sequence of this 7 kDa wheat nsLtp protein shows a high degree of similarity with the predicted protein from the open reading frame (ORF) of the Bz11E cDNA,
10 which had been isolated in a differential screening for barley aleurone specific transcripts (Jakobsen *et al.*, 1989). However, the amino acid sequence of this polypeptide showed only limited sequence identities with the previously sequenced 10 kDa proteins. In sub-cellular localisation studies using gold labelled antibodies one 10 kDa protein from *Arabidopsis* was localised to the cell wall of epidermal leaf cells.
15 The presence of a signal peptide domain in the N-terminus of the open reading frames of all characterised plant nsLtp cDNAs, also suggests that these are proteins destined for the secretory pathway with a possible extracellular function.

Olsen *et al.* in a paper titled "Molecular Strategies For Improving Pre-Harvest
20 Sprouting Resistance In Cereals" published in 1990 in the published extracts from the Fifth International Symposium On Pre-Harvest Sprouting In Cereals (Westview Press Inc.) describe three different strategies for expressing different "effector" genes in the aleurone layer in developing grains of transgenic plants. This document mentions 4 promoter systems - including a system called B11E.

25

Kalla *et al.* (1993) in a paper titled "Characterisation of Promoter Elements Of
Aleurone Specific Genes From Barley" describe the possibility of the expression of anti-sense genes by the use of promoters of the aleurone genes B22E, B23D, B14D, and B11E.

30

Linnestad *et al.* (1991) describe the isolation and sequencing of the Ltp1 gene and disclose a 787 base pair fragment of the Ltp1 gene promoter fused to a fragment of the Ltp1 structural gene. This paper does not disclose any expression studies using the 787 base pair fragment.

5

Skriver *et al.* (1992) report further on the Ltp1 gene. This paper says that the Ltp1 gene promoter is only aleurone specific. To confirm this submission the paper further reports on the isolation and fusion of a 769 bp fragment (-702 to +67 bp) of the gene to the bacterial β -glucuronidase (*GUS*) reporter gene. This fragment
10 therefore contains 635 bp of the Ltp1 gene promoter. Subsequent transient expression studies showed that the shortened gene promoter resulted only in aleurone specific expression. Expression was not observed in any other tissue. The authors conclude that there are sequences between the -702 and +67 bp of Ltp1 which contain DNA elements that specifically modulate its transcription in aleurone cells.

15

One of the major limitations to the molecular breeding of new types of crop plants with specific cells expressing GOIs is the lack of a suitable tissue specific promoter. In particular, there is a lack of a tissue specific promoter that leads to expression of a GOI in a developing caryopsis (or grain) or in a germinating rice seedling or in a
20 developing grain, in particular in the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant.

Moreover, all of the available promoters - such as the CaMV 35S, rice actin and maize alcohol dehydrogenase - are constitutive, i.e. they are fairly non-specific in
25 target site or stage development as they drive expression in most cell types in the plants.

Hence, another problem that arises is how to achieve expression of a product coded for by a GOI in a specific tissue that gives minimal interference with the developing
30 embryo and seedling.

Our co-pending United Kingdom patent application (GB 9324707.0) describes the use of an Ltp2 gene promoter for expression of a GOI in the aleurone layer. However, in spite of this teaching, there is still a need for other tissue specific promoters, such as another aleurone specific promoter or, preferably, a promoter specific for vascular tissue and/or the scutellar epithelial layer. In this regard, it is still desirable to provide other tissue specific expression of GOIs in cereals such as rice, maize, wheat, barley and other transgenic cereal plants. Moreover it is desirable to provide tissue specific expression that does not detrimentally affect the developing embryo and the developing caryopsis (or grain).

10

According to a first aspect of the present invention there is provided a modified Ltp1 gene promoter which is integrated, preferably stably integrated, within a plant material's genomic DNA and which is capable of inducing expression of a GOI when fused to the gene promoter in at least the aleurone cells or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant (e.g. in the root, leaves and stem).

15

According to a second aspect of the present invention there is provided a modified Ltp1 gene promoter according to claim 1 or claim 2 wherein the promoter comprises the nucleic acid sequence shown as SEQ. I.D. 1, or a sequence that has substantial homology with that of SEQ. I.D. 1, or a variant thereof.

20

According to a third aspect of the present invention there is provided an isolated Ltp1 gene promoter comprising the sequence shown as SEQ. I.D. 1, or a sequence that has substantial homology therewith, or a variant thereof.

25

According to a fourth aspect of the present invention there is provided a construct comprising a GOI and a modified Ltp1 gene promoter according to the present invention; wherein the construct is capable of being expressed in at least the aleurone cells or in at least the scutellar epithelial tissue or vascular tissue of a plant material; and wherein if there is expression in just the aleurone layer of a developing barley caryopsis then the fused promoter and GOI are not the 769 bp fragment of Skriver

30

et al (1992).

According to a fifth aspect of the present invention there is provided an expression system for at least the aleurone cells or for at least the scutellar epithelial tissue or
5 vascular tissue of a plant material, the expression system comprising a GOI fused to a modified Ltp1 gene promoter wherein the expression system is capable of being expressed in at least the aleurone cells or in at least the scutellar epithelial tissue or vascular tissue of the plant material; and wherein if there is expression in just the aleurone layer of a developing barley caryopsis then the fused promoter and GOI are
10 not the 769 bp fragment of Skriver *et al* (1992).

According to a sixth aspect of the present invention there is provided an expression system for at least the aleurone cells of a developing caryopsis or for at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or developing
15 grain or plant (e.g. in the root, leaves and stem), the expression system comprising a gene promoter fused to a GOI wherein the expression system is capable of being expressed in at least the aleurone cells of the developing caryopsis or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant (e.g. in the root, leaves and stem); either wherein if there is
20 expression in just the aleurone layer of a developing barley caryopsis then either the promoter is not the wild type Ltp1 promoter in its natural environment and the GOI is not the Ltp1 functional gene in its natural environment; or wherein if there is expression in just the aleurone layer of a developing caryopsis then the fused promoter and GOI are not the 769 bp fragment of Skriver *et al* (1992).

25

According to a seventh aspect of the present invention there is provided a transgenic cereal comprising an expression system according to the present invention or a construct according to the present invention wherein the expression system or construct is integrated, preferably stably integrated, within the cereal's genomic
30 DNA.

According to an eighth aspect of the present invention there is provided the use of a gene promoter according to the present invention to induce expression of a GOI when fused to the gene promoter in at least the aleurone cells or in at least the scutellar epithelial tissue or vascular tissue of a plant material.

5

According to a ninth aspect of the present invention there is provided a process of expressing a GOI when fused to a gene promoter according to the present invention wherein expression occurs in at least the aleurone cells or in at least the scutellar epithelial tissue or vascular tissue of a plant material.

10

According to a tenth aspect of the present invention there is provided a process of expressing in at least the scutellar epithelial tissue or vascular tissue of a developing grain or a germinating seedling or a plant, preferably a developing rice grain or a germinating rice seedling or a transgenic rice plant, an expression system according to the present invention or a construct according to the present invention wherein the expression system or construct is integrated, preferably stably integrated, within the cereal's genomic DNA.

15

According to an eleventh aspect of the present invention there is provided a combination expression system comprising a. as a first construct, a construct according to the present invention; and b. as a second construct, a construct comprising a GOI and another gene promoter that is tissue- or stage-specific.

20

According to a twelfth aspect of the present invention there is provided a developing cereal grain, preferably a germinating rice seedling, comprising any one of: a promoter according to the present invention, an expression system according to the present invention, a construct according to the present invention, or a combination expression system according to the present invention.

25

According to a thirteenth aspect of the present invention there is provided plasmid NCIMB 40609.

30

Preferably the plant material is a developing caryopsis, a germinating seedling, a developing grain or a plant.

5 Preferably the construct is capable of being expressed in at least the aleurone cells of a developing caryopsis or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant when the construct is integrated, preferably stably integrated, within the caryopsis's or grain's or seedling's or plant's genomic DNA.

10 Preferably the modified Ltp1 gene promoter comprises the nucleic acid sequence shown as SEQ. I.D. 1, or a sequence that has substantial homology with that of SEQ. I.D. 1, or a variant thereof.

15 Preferably the construct further comprises at least one additional sequence to increase expression of the GOI.

20 Preferably the expression system is for at least the aleurone cells of a developing caryopsis or for at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or developing grain or plant (e.g. in the root, leaves and stem).

Preferably the expression system is additionally capable of being expressed in the embryo cells of the germinating grain or the plantlet.

25 Preferably the expression system is integrated, preferably stably integrated, within a developing caryopsis's genomic DNA or a germinating seedling's genomic DNA or a developing grain's genomic DNA or a plant's genomic DNA.

30 Preferably, in the expression system, the gene promoter comprises the sequence shown as SEQ I.D. No. 1 or comprises a sequence that has substantial homology therewith, or is a variant thereof.

Preferably, the expression system comprises the construct according to the present invention.

5 Preferably, in the use, the gene promoter is used to induce expression of a GOI when fused to the gene promoter in at least the aleurone cells of a developing caryopsis or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant (e.g. in the root, leaves and stem).

10 Preferably, the gene promoter expresses the GOI when fused to the gene promoter in at least the aleurone cells of a developing caryopsis or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant (e.g. in the root, leaves and stem).

15 Preferably the promoter and GOI are integrated, preferably stably integrated, within a cereal's genomic DNA.

Preferably the gene promoter is a fragment of a barley Ltp1 gene promoter.

20 Preferably the promoter is for a 10 kDa lipid transfer protein.

Preferably the gene promoter is obtainable from plasmid NCIMB 40609.

25 Preferably the gene promoter is used for expression of a GOI in a cereal caryopsis or a cereal grain or a cereal seedling or a cereal plant.

Preferably the cereal caryopsis is a developing cereal caryopsis, the cereal grain is a developing cereal grain, and the cereal seedling is a germinating cereal seedling.

30 Preferably the cereal is any one of a rice, maize, wheat, or barley.

Preferably the cereal is rice or maize.

Preferably the developing caryopsis is a developing barley caryopsis, the germinating seedling is a germinating rice seedling, the developing grain is a developing rice grain, and the plant is a transgenic rice plant.

- 5 Preferably in the combination expression system each construct is integrated, preferably stably integrated, within a plant material.

Preferably each of the myb site and the myc site in the gene promoter is maintained substantially intact.

10

- Preferably the gene promoter is integrated, preferably stably integrated, in the developing caryopsis's genomic DNA or the germinating seedling's genomic DNA or the developing grain's genomic DNA or the plant's genomic DNA and which is capable of inducing expression of a GOI when fused to the gene promoter in at least
15 the aleurone cells of the developing caryopsis or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant (e.g. in the root, leaves and stem).

- Preferably the transgenic developing caryopsis, germinating seedling, developing
20 grain or plant is prepared by stable integration of the GOI and the gene promoter to form a stable transgenic plant. This ensures aleurone or epithelial or vascular expression at, at least, the developing caryopsis stage. One preferred method for achieving this includes preparing the transgenic developing caryopsis, germinating seedling, developing grain or plant by stable integration of the GOI and the gene
25 promoter at the protoplast level.

- Preferably the promoter is used for expression of a GOI in a monocotyledonous species, including a grass - preferably a transgenic cereal grain or caryopsis. Preferably the gene promoter is used for expression of a GOI in a cereal grain or
30 caryopsis. Preferably the cereal grain or caryopsis is a developing cereal grain or caryopsis. Preferably the cereal grain or caryopsis is any one of a rice, maize, wheat, or barley grain or caryopsis.

13

Preferably the cereal grain is a rice grain.

Preferably the DNA sequence for the modified Ltp1 gene promoter is the nucleic acid sequence shown as SEQ. I.D. 1.

5

Preferably in the combination expression system each construct is integrated, preferably stably integrated, within a developing caryopsis's genomic DNA or a grain's genomic DNA or a seedling's genomic DNA or a plant's genomic DNA.

- 10 Preferably, in the combination expression system, the first construct comprises the modified Ltp1 gene promoter according to the present invention.

Preferably, the promoter in the second construct is an aleurone specific promoter.

- 15 Preferably the promoter in the second construct is a barley promoter.

Preferably the second construct is the B22E gene promoter.

Preferably the promoter in the second construct is the Ltp2 gene promoter.

20

Preferably the promoter in the second construct is for a 7 kDa lipid transfer protein.

Preferably the promoter in the second construct is the promoter for Ltp2 of *Hordeum vulgare*.

25

Preferably the promoter in the second construct comprises the sequence shown as SEQ. I.D. 2, or a sequence that has substantial homology therewith, or a variant thereof.

- 30 Preferably each of the myb site and the myc site in the Ltp2 gene promoter is maintained substantially intact.

Preferably the second construct further comprises at least one additional sequence to increase expression of the GOI.

Preferably, in the combination expression system, the grain or caryopsis is as defined
5 above for the present invention.

Preferably the gene promoter is obtainable from plasmid NCIMB 40609.

A preferred embodiment of the present invention is a modified Ltp1 gene promoter
10 which is integrated, preferably stably integrated, within a plant material's genomic DNA and which is capable of inducing expression of a GOI when fused to the gene promoter in at least the aleurone cells or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant (e.g. in the root, leaves and stem), but wherein if there is expression in just the aleurone layer
15 of a developing seed then the fused promoter and GOI are not the 769 bp fragment of Skriver *et al* (1992).

An even more preferred embodiment of the present invention is a modified Ltp1 gene promoter which is integrated, preferably stably integrated, within a plant material's
20 genomic DNA and which is capable of inducing expression of a GOI when fused to the gene promoter in at least the aleurone cells or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant (e.g. in the root, leaves and stem), wherein the promoter comprises the nucleic acid sequence shown as SEQ. I.D. 1, or a sequence that has substantial homology with
25 that of SEQ. I.D. 1, or a variant thereof.

As a highly preferred embodiment, the present invention therefore provides transgenic rice comprising a construct comprising a GOI fused to a modified Ltp1 gene promoter; wherein the construct is integrated, preferably stably integrated, within the
30 rice's genomic DNA, and wherein the GOI is expressed in at least the vascular tissue and/or scutellar epithelial layer of a germinating rice seedling or a developing rice grain or a rice plant.

In a more preferred embodiment the present invention provides a transgenic rice seedling, grain or plant comprising a construct comprising a GOI fused to a modified Ltp1 gene promoter, wherein the construct is integrated, preferably stably integrated, within the rice's genomic DNA; wherein the GOI is expressed in at least the scutellar
5 epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant, and wherein the modified Ltp1 gene promoter comprises the nucleic acid sequence shown as SEQ. I.D. 1, or a sequence that has substantial homology with that of SEQ. I.D. 1, or a variant thereof.

10 The additional sequence(s) for the construct(s) for increasing the expression of the GOI(s) may be one or more repeats (e.g. tandem repeats) of the promoter upstream box(es) which are responsible for the aleurone layer or scutellar epithelial cell and/or vascular expression pattern of the modified Ltp1 gene promoter. The additional sequence may even be a *ShI*-intron.

15

The term "plant material" includes a developing caryopsis, a germinating caryopsis or grain, or a seedling, a plantlet or a plant, or tissues or cells thereof, such as the aleurone cells of a developing caryopsis or the scutellar epithelial tissue or vascular tissue of a germinating seedling or developing grain or plant (e.g. in the root, leaves
20 and stem).

Thus a preferred aspect of the present invention comprises plant material comprising a GOI and a modified Ltp1 gene promoter which is capable of inducing expression of the GOI when fused to the gene promoter in at least the aleurone cells or in at least
25 the scutellar epithelial tissue or vascular tissue of the plant material; wherein the construct is capable of being expressed in at least the aleurone cells or in at least the scutellar epithelial tissue or vascular tissue of the plant material, when the construct is integrated, preferably stably integrated, within the caryopsis's or grain's or seedling's or plant's genomic DNA; and wherein the modified Ltp1 gene promoter
30 comprises the nucleic acid sequence shown as SEQ. I.D. 1, or a sequence that has substantial homology with that of SEQ. I.D. 1, or a variant thereof.

The term "modified" with reference to the present invention means any Ltp1 gene promoter that is different to the wild type promoter but wherein the promoter induces expression in at least the aleurone cells of a developing caryopsis or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant (e.g. in the root, leaves and stem).

In particular, a preferred modified Ltp1 gene promoter is a shortened wild type Ltp1 gene promoter but wherein the promoter induces expression in at least the aleurone cells of a developing caryopsis or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant (e.g. in the root, leaves and stem).

The term "transgenic" in relation to the present invention - in particular in relation to the developing caryopsis, germinating seedlings, developing grains and plants of the present invention - does not include a wild type promoter in its natural environment in combination with its associated functional gene (GOI) in its natural environment. Thus, the term includes developing caryopsis or seedlings or grains or plants incorporating the GOI which may be natural or non-natural to the grain or caryopsis or seedling or grain or plant in question operatively linked to the modified Ltp1 promoter of the present invention.

The term "GOI" with reference to the present invention means any gene of interest. A GOI can be any gene that is either foreign or natural to the cereal in question, except for the wild type Ltp1 functional gene when in its natural environment. In the combination expression system the GOI may be the same or different.

Typical examples of a GOI include genes encoding for proteins and enzymes that modify metabolic and catabolic processes. For example, the GOI may be a protein giving added nutritional value to the grain or caryopsis as a food or crop. Typical examples include plant proteins that can inhibit the formation of anti-nutritive factors and plant proteins that have a more desirable amino acid composition (e.g. a higher lysine content than the non-transgenic plant).

The GOI may even code for an enzyme that can be used in food processing such as chymosin, thaumatin, α -galactosidase and guar.

In a preferred embodiment, particularly with vascular expression, the GOI may code
5 for an agent for introducing or increasing pathogen resistance.

The GOI may even be an antisense construct for modifying the expression of natural transcripts present in the relevant tissues.

10 The GOI may even code for a non-natural plant compound that is of benefit to animals or humans. For example, the GOI could code for a pharmaceutically active protein or enzyme such as the therapeutic compounds insulin, interferon, human serum albumin, human growth factor and blood clotting factors. In this regard, the transformed cereal grain or caryopsis could prepare acceptable quantities of the
15 desired compound which could be easily retrievable from the scutellar epithelial layer, the aleurone layer or the vascular tissue.

Preferably the GOI is a gene encoding for any one of a protein having a high nutritional value, a *Bacillus thuringensis* insect toxin, an α - or β - amylase antisense
20 transcript, a protease antisense transcript, or a glucanase antisense transcript.

The term "a variant thereof" with reference to the present invention means any substitution of, variation of, modification of, replacement of, deletion of or the addition of one or more nucleic acid(s) from or to the promoter sequence providing
25 the resultant sequence exhibits at least aleurone, scutellar epithelial or vascular expression, respectively. The term also includes sequences that can substantially hybridise to the promoter sequence.

The term "substantial homology" covers homology with respect to at least the
30 essential nucleic acids/nucleic acid residues of the promoter sequence providing the homologous sequence acts as a promoter, e.g. as a promoter for at least aleurone expression in a developing caryopsis or for at least scutellar epithelial tissue or

vascular tissue expression in a germinating seedling or in a developing grain or plant. Preferably there is at least about 80% homology, more preferably at least about 90% homology, and even more preferably there is at least about 95% homology with the promoter sequence shown as SEQ. I.D. No. 1. or SEQ. I.D. No. 2, respectively.

5

The term "maintained substantially intact" means that at least the essential components of each of the myb site and the myc site remain in the construct to ensure acceptable expression of a GOI. Preferably at least about 75%, more preferably at least about 90%, and even more preferably there is at least about 95%, of the myb or myc site is left intact.

10

The term "construct" - which is synonymous with terms such as "cassette", "hybrid" and "conjugate" - includes a GOI directly or indirectly attached to the modified gene promoter, such as to form a [modified Ltp1 gene promoter-GOI] cassette. An example of an indirect attachment is the provision of a suitable spacer group such as an intron sequence, such as the *Shl*-intron, intermediate the promoter and the GOI. The same is true for the term "fused" in relation to the present invention which includes direct or indirect attachment.

15

20 The term "expression system" means that the system defined above can be expressed in an appropriate organism, tissue, cell or medium. In this regard, the expression system of the present invention may comprise additional components that ensure to increase the expression of the GOI by use of the gene promoter.

25 As indicated above, the expression system of the present invention can also be used in conjunction with another expression system, preferably an expression system that is also tissue and/or stage specific.

For example, the construct comprising the modified Ltp1 gene promoter (e.g. the 787 bp fragment of SEQ. I.D. NO. 1) can be used in conjunction with a construct comprising the Ltp2 gene promoter (e.g. SEQ. I.D. NO. 2) - which is the subject of our co-pending UK patent application GB 9324707.0.

30

In this respect, and with reference to barley, in the early stages of developing caryopsis the modified Ltp1 gene promoter affects expression of a GOI in at least the aleurone layers of developing caryopsis. This expression can then be complimented by use of the Ltp2 gene which can express a GOI (which may be the same or different as that operatively linked to the modified Ltp1 gene promoter) in high levels in the aleurone layer of developing grains.

However, the combination expression system is very effective for transgenic rice. In this respect, in the early stages of developing caryopsis the modified Ltp1 gene promoter expresses a GOI in the scutellar epithelial layer and the vascular tissue. This expression can then be complimented by use of the Ltp2 gene which can express a GOI in high levels in the aleurone layer of developing grains. This combination is particularly advantageous for pre-harvest sprouting when the first response is production of α -amylase in the scutellar epithelium cells as this can be reduced or prevented by placing an anti-sense α amylase gene under the control of the Ltp1 promoter. In this system, the expression of antisense α -amylase would block the synthesis of α -amylases in the scutellum epithelial cells - where they are first made. The same or another GOI could be expressed in the aleurone layer via the Ltp2 gene promoter.

20

The construct comprising the modified Ltp1 gene promoter may even be used in conjunction with a construct comprising the B22E gene promoter - details of which may be found in Olsen *et al.* (1990) and Klemsdal *et al.*, (1991). This gene promoter, which is expressed in immature aleurone layers, has been shown by particle bombardment experiments to be capable of driving *Gus* expression in developing barley grains. Also, using Northern analysis, as well as *in situ* hybridization, it has been shown that the B22E cDNA probe hybridizes to transcripts in the aleurone layer and in the scutellum parenchyma cells and the provascular bundle of the embryo axis in developing barley grains. In addition, a hybridizing transcript is also present in the ventral vascular strand of developing caryopsis (Olsen *et al.*, 1990).

30

We have also found that by using a 4.6 kb B22E promoter fragment contained on a XbaI-ClaI fragment of a genomic clone fused to the *Gus* reporter gene transformed rice plants could be prepared. Those transformed rice plants exhibited strong expression in the vascular tissue (phloem) of the ventral strand of the developing rice grain. This expression pattern was completely unexpected in view of Klemsdal *et al* (1991). Expression, although weaker, in the same cell type was also observable in the stem of young shoots. Thus, using the B22E promoter, a GOI transcript can be expressed in the aleurone layers of developing grains, the parenchyma cells of the embryonic scutellum and the ventral vascular bundle of developing grains.

10

The combination of the use of the modified Ltp1 gene promoter and the B22E gene promoter could even include the use of another gene promoter, such as the Ltp2 gene promoter, to express three GOIs respectively wherein each GOI may be the same or different.

15

One or more of the other expression systems to be used in conjunction with the modified Ltp1 gene promoter expression system may be contained in or on the same transmission vector - such as in the same transforming bacterium or even in the same plasmid. The advantage of this is that each expression system can then be delivered at the same time. The respective expression systems will then be turned on during the relevant life time of the grain or caryopsis or the plantlet or the mature plant.

20

The present invention therefore provides the novel and inventive use of a promoter which can express a GOI in at least the aleurone cells of a developing caryopsis or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant (e.g. in the root, leaves and stem). In a preferred embodiment the present invention relates to the use of a modified Ltp1 gene promoter, preferably the Ltp1 gene promoter is obtainable from barley.

25

The main advantage of the present invention is that the use of the modified Ltp1 gene promoter results in expression of a GOI in at least the aleurone layer of at least a developing caryopsis, such as a developing barley caryopsis, or in at least the

30

scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or plant of cereals such as rice, maize, wheat or other transgenic cereal grain or caryopsis, preferably a developing rice grain.

- 5 Another advantage is that, depending on the type of GOI, the expressed products can be stable *in vivo*. Hence over a period of time high levels of the expressed product can accumulate in the aleurone cells or in epithelial cells or in the vascular tissue.

10 A further advantage is that the expression of the product coded for by a GOI in the aleurone layer or the epithelial layer or the vascular tissue has minimal interference with the developing embryo and seedling. This is in direct contrast to known constitutive promoters which give high levels of expression in the developing seedling and mature plant tissues which severely affect normal plant development. Thus the present invention is particularly useful for expressing a GOI in at least the aleurone
15 layer of a developing caryopsis or in at least the scutellar epithelial tissue or in the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or plant - such as cereal grains or caryopsis - and in doing so not detrimentally affect the caryopsis, seedling, grain or plant.

- 20 With regard to the first aspect of the present invention it is to be noted that this is the first reported case for the specific expression of a GOI in the scutellar epithelial cells or vascular cells of a transformed developing cereal grain such as rice.

25 With regard to some aspects of the present invention, it is to be noted that up until now it was believed that the wild type Ltp1 gene promoter or a specific variant thereof when fused to at least a segment of the Ltp1 functional gene would lead only to expression in the aleurone layer. For example see the teachings of Skriver *et al.* (1992). However, with the present invention, we have now surprisingly found that this is not the case and it is now possible to modify the Ltp1 gene promoter to lead
30 to a pronounced expression in at least the aleurone layer or in at least the scutellar epithelial layer or vascular tissue of a plant material.

In one embodiment the plant material is barley plant material. In another embodiment the plant material is not barley plant material. In a preferred embodiment the plant material is rice plant material. In an alternative preferred embodiment the plant material is maize plant material.

5

In a germinating, transgenic barley caryopsis according to the present invention, there is expression in the aleurone layer.

10 In a germinating, transgenic rice seedling according to the present invention there is pronounced expression in the scutellar epithelial tissue and vascular tissue.

As indicated, the expression pattern for the present invention is particularly surprising as it was completely unexpected that a modified Ltp1 gene promoter could result in expression of a GOI, such as a plant functional gene, in the aleurone cells of, for example, barley or in the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or plant of rice (see experimental section later). The findings of the present invention are also surprisingly different to the work of Skriver *et al.* (1992) who, as mentioned above, report that the Ltp1 gene promoter and a shortened version thereof when fused to the functional Ltp1 functional gene only result in aleurone specific expression in barley - i.e. expression is not observed in any other tissue in barley or even other cereals.

20

In order to prepare the transgenic organism according to the present invention, the modified Ltp1 gene promoter may be initially inserted into a plasmid. For example, the *SacI-BcII* Ltp1 gene promoter fragment can be inserted into the *SacI-BamHI* site of Bluescript. A GOI, such as *GUS*, can then be inserted into this construct. Furthermore, a *Sh1* intron can then be inserted into the *SmaI* site of this construct.

25

Stable integration into protoplasts may be achieved by using the method of Shimamoto (1989). Another way is by bombardment of an embryonic suspension of cells (e.g. rice, barley or maize cells). A further way is by bombardment of immature embryos (e.g. rice, maize or barley embryos).

30

With regard to the present invention, it is shown by using particle bombardments that the modified *Ltp1* gene promoter, such as the 787 bp fragment of the attached sequence, when fused to a β -glucuronidase (*GUS*) reporter gene, which serves as a GOI for the purposes of this invention, acts as a promoter for expression of *GUS* in a specific tissue type or specific tissue types. For example, *GUS* expression can be achieved in the aleurone cells of developing cereal caryopsis or grain, in particular developing barley caryopsis, or in the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or plant, in particular developing rice grain or germinating seedlings.

10

In particular, in transgenic rice plants, the modified barley *Ltp1* gene promoter directs strong expression of the *GUS*-reporter gene in the scutellar epithelial layer and the vascular tissue of the developing caryopsis. This expression can continue through into the germinating grains. The surprising finding is that very pronounced expression can be seen in the scutellar epithelial tissue or vascular tissue of a developing rice grain or germinating rice seedlings. Other examples include expression in the vascular bundles and tip of emerging shoots and roots, leaf veins and vascular bundles of stems.

15

Generally therefore the present invention relates to a modified promoter for a *Ltp1* gene encoding a 10 kDa nsLTP. In the present invention, a genomic clone was isolated using the cDNA insert of previously isolated cDNA clone and characterised by DNA sequencing (see discussion later). The sequence of the cDNA and isolated genomic clone was found to be identical in the overlapping region. It was found the *Ltp1* gene contains one intron (see discussion later).

20
25

By comparing the DNA sequence of the active promoter sequences two putative *cis*-acting elements with the potential of binding known transcriptional factors present in cereals were detected. They include the binding sites for transcriptional factors of the myb and myc class, namely TAACTG and CANNTG respectively. Our studies showed that high levels of expression are achieved when the myb and myc sites are left intact.

30

In the present invention, mature fertile rice plants were regenerated from transformed cultured rice protoplasts. The developing caryopsis of these primary transformants were analysed for the expression of *GUS*. It was found that the modified barley *Ltp1* gene promoter confers some expression in the aleurone layer of the transgenic rice plants. However, pronounced expression was observed in the scutellar epithelial tissue or vascular tissue of germinating rice seedlings or developing transgenic rice grain or transgenic rice plants. This is the first example of such patterns of expression in transgenic rice plants.

The following sample has been deposited in accordance with the Budapest Treaty at the recognised depositary The National Collections of Industrial and Marine Bacteria Limited (NCIMB) at 23 St Machar Drive, Aberdeen, Scotland, AB2 1RY, United Kingdom, on 11 January 1994:

An *E. Coli* K12 bacterial stock containing the plasmid pLtp1.787-GN - i.e. Bluescript containing a 787 bp fragment of the barley *Ltp1* gene promoter (Deposit Number NCIMB 40609).

The plasmid pLtp1.787-GN is shown pictorially in Figure 6 (see later).

The modified *Ltp1* gene promoter can be isolated from this plasmid through the use of appropriate PCR primers, which may be easily constructed from the data from the shown sequences.

Other embodiments and aspects of the present invention include:

A transformed host having the capability of expressing a GOI in the aleurone layer or the scutellar epithelial layer or the vascular tissue through the use of the gene promoter as hereinbefore described;

A vector incorporating a construct as hereinbefore described or any part thereof;

25

A plasmid comprising a construct as hereinbefore described or any part thereof;

5

A cellular organism or cell line transformed with such a vector;

A monocotyledonous plant comprising any one of the same;

A developing caryopsis or grain or germinating seedling comprising any of the same; and

10

A method of expressing any one of the above.

The present invention will now be described only by way of examples in which reference shall be made to the accompanying Figures in which:

15

Figure 1 is a diagrammatic illustration of the structural components of a developing caryopsis;

20

Figure 2 shows the results for an *in situ* hybridization experiment for a wild type Ltp1 gene promoter in barley;

Figure 3 is a nucleotide sequence of part of the wild type Ltp1 gene taken from Linnestad *et al.* (1991);

25

Figure 4 is a nucleotide sequence of part of the wild type Ltp1 gene taken from Skriver *et al.* (1992);

Figure 5 is a nucleotide sequence of a 787 bp fragment of the wild type Ltp1 gene promoter;

30

Figure 6 is a linear map of the Ltp1.787-GN construct showing additional sequence information;

Figure 7 is a circular map of the plasmid pLtp1.787-GN containing the Ltp1.787-GN construct;

5 Figure 8 is a longitudinal section of a developing rice grain post expression of the modified Ltp1 gene promoter; and

Figure 9 is a longitudinal section of a mature germinating rice grain post expression of the modified Ltp1 gene promoter.

10 A. METHODS

i. Plant material

15 Caryopsis of barley (*Hordeum vulgare* cv. Bomi) were collected from plants grown in a phytotron as described before (Kvaale and Olsen, 1986). The plants were emasculated and pollinated by hand and isolated in order to ensure accurate determination of caryopsis age.

ii. cDNA and genomic clones

20

The isolation and sequencing of the Ltp1 cDNA clone was conducted as described by Jakobsen *et al.* (1989). A barley, cv. Bomi genomic library was constructed by partial *Mbo*I digestion of total genomic DNA and subsequent ligation of the 10-20 kilo basepair (kb) size fraction with *Bam*HI digested lambda EMBL3 DNA (Clontech
25 Labs, Palo Alto, Ca, USA). Using the Ltp1 cDNA insert as a template for probe synthesis with a random labelling kit (Boehringer-Mannheim), one positive clone was identified after repeated rounds of plaque hybridization. DNA purified from this clone was restricted with several enzymes and characterized by Southern blot analysis. The sequence data obtained after this procedure are shown in Figure 3.

30

iii. *In situ* hybridization

For *in vitro* transcription of antisense RNA, the plasmid Ltp1 was linearized and transcribed using MAXIscript (Ambion) and [α 32 P]-UTP (Amersham International).

- 5 The probe was hydrolysed to fragments of about 100 bp as described by Somssich *et al.* (1988). Caryopsis tissues were fixed in 1% glutaraldehyde, 100 mM sodium phosphate (pH 7.0) for 2 hours and embedded in Histowax (Histolab, Göteborg, Sweden).

- 10 Barley caryopsis sections of 10 μ m were pre-treated with pronase (Calbiochem) as described by (Schmelzer *et al.*, 1988) and hybridized with 25 ml of hybridization mix (200 ng probe ml⁻¹, 50% formamide, 10% (w/v) dextran sulphate, 0.3 M NaCl, 10 mM Tris-HCl, 1 mM EDTA (pH 7), 0.02% polyvinyl-pyrrolidone, 0.2% Ficoll, 0.02% bovine serum albumin) for 15 hours at 50 °C.

15

Post-hybridization was carried out according to Somssich *et al.* (1988) and autoradiography was done as described by Schmelzer *et al.* (1988).

iv. Constructs for transient expression analysis

20

For the micro-projectile bombardment experiments, the following was used:

pLtp1.787-GN (see Figure 7 and associated commentary).

- 25 Isolated plasmid DNA was used in the bombardment studies.

For transient assay studies with rice protoplasts, the following were studied:

pLtp1.787-GN (see Figure 7 and associated commentary).

30

pLtp1.787(-myb/myc)-GN (see commentary below).

Deletion studies were performed on the modified Ltp1 gene promoter (Ltp1.787) wherein a section of DNA containing the myb and myc sites (see Figure 3 and associated commentary) was removed to form pLtp1.787(-myb/myc)-GN. In this embodiment, the modified Ltp1 gene promoter having deletions from and between the myb and myc sites was prepared and fused to GN. In order to prepare this deleted modified Ltp1 gene promoter a PCR strategy using primers covering the flanking sequences of the deleted sequence was adopted.

v. Transformation of barley cells by particle bombardment

10

Barley caryopsis were harvested at 25 DAP (days after pollination), surface sterilized in 1% sodium hypochlorite for 5 min and then washed 4 times in sterile distilled water. The maternal tissues were removed to expose the aleurone layer and the caryopsis was then divided into two, longitudinally along the crease. The pieces of tissue were then placed, endosperm down, onto MS media (Murashige & Skoog 1962) with 10 g/l sucrose solidified with 10 g/l agar in plastic petri dishes (in two rows of 4 endosperm halves per dish). Embryos from the same caryopsis were placed in the same petri dishes with the scutellum side facing upwards.

15

Single bombardments were performed in a DuPont PDS 1000 device, with M-17 tungsten pellets (approx. 1 μ m in diameter) coated with DNA as described by Gordon-Kamm *et al.* (1990) and using a 100 mm mesh 2 cm below the stopping plate. Histochemical staining for *GUS* expression was performed with X-Gluc (5-bromo,4-chloro,3-indolyl, β -D-Glucuronic acid) as described by Jefferson (1987) at 37°C for 2 days.

20

In these studies, after bombardment with the pLtp1.787-GN and staining for *GUS*-activity, blue spots appeared both in the aleurone layer as well as in the scutellar epithelium layer. These results demonstrate that the 787 bp fragment of the Ltp1 gene promoter of the present invention is capable of driving transcription in the epithelial cells.

25

30

vi. Rice transformation

In these studies, the gene was transformed into rice by electroporation of embryogenetic protoplasts following the teachings of Shimamoto et al. 1989. Six
5 fertile transgenic rice plants were obtained. Histochemical *GUS* analysis was also carried out with developing rice grains of 25 DAP and 1 to 5 day old seedlings and up to 1 month old plants derived from transgenic grains. The results demonstrated expression of the Ltp1 - *GUS* gene in the scutellar epithelial layer of developing transgenic rice plants. In addition, in a germinating rice seedling according to the
10 present invention there is a pronounced expression in the vascular tissue.

B. RESULTS AND DISCUSSION WITH REFERENCE TO THE FIGURES

1. In order to explain more fully the results, reference is made to Figure 1 which
15 shows the major components of a typical developing caryopsis (or grain) 1. In this regard, the caryopsis (or grain) 1 comprises an endosperm component 3 and an embryo component 5. The endosperm component 3 is divisible into an outer aleurone layer 7, which is three cells thick for barley caryopsis, and a starchy endosperm 9. The embryo component 5 is divisible into a scutellum 11 and an embryo axis 13.
20 The scutellum 11 is further divisible into an epithelial layer 15 and parenchyma layer 17. Likewise, the embryo axis 13 is further divisible into a root component 19 and a shoot component 21.

2. Figure 2 is a transverse section of a 30 day-old wild-type developing barley
25 caryopsis showing *in situ* hybridisation with a radio-labelled Ltp1 probe. The bound probe is only seen in the aleurone layer. It is not seen in any other tissue type, in particular the scutellar epithelial layer. This work confirms the work of Skriver *et al.* (1992).

30 The bright spots are due to optical interference.

3. Figure 3 shows the nucleotide sequence and the deduced amino acid sequence of Ltp1. The intron is indicated by lower case letters. The TGA stop codon is indicated by an asterisk, the putative CAAT and TATA sequences are indicated by boxes. A 21 bp inverted repeat is indicated by arrows. Four 8 bp palindromic sequences are overlined. The motif indicated by thick underlining resembles the CATGTAAA motif present in the promoters of several genes expressed in aleurone cells (Klemsdal *et al.* (1991)). An AT block followed by a myb consensus recognition site and a myc binding motif are indicated by double underlining.
4. Figure 4 shows the sequence of the Ltp1 gene. The 351 bp open reading frame is interrupted by a 133 bp intron (+412 to +544). The transcript start site is at position +1. The putative CAAT and TATA boxes are at -107 and -34. A putative poly (dA) site is at +785 (Skriver *et al.* (1992)).
5. Figure 5 is the nucleotide sequence of the preferred embodiment of the present invention, i.e. a 787 bp fragment of the Ltp1 gene promoter. The same commentary for Figure 3 is equally applicable here.
6. Figure 6 is an outline of the Ltp1 genomic clone containing the Ltp1 structural gene (shaded box) and the promoter fragment fused to the *GUS* gene (black box) used to transform rice. Also indicated are the extensions of the Ltp1 fragment described in Linnestad *et al.* (1991) and Skriver *et al.* (1992). The figures used represent DNA fragment lengths in kb. The total length of the genomic clone is in the order of 8.1 kb.
7. Figure 7 helps explain how pLtp1.787-GN was constructed. In this regard, the following fragments were sequentially cloned into the vector Bluescript KS: firstly the 787 bp SacI/Bcl fragment of the Ltp1 gene promoter was cloned into the SacI/BclII site of the vector; and secondly a *GUS*-Nos Terminator on 2150 bp SmaI/EcoRI fragment derived from pBI101 was cloned into SmaI/EcoRI downstream of the Ltp1 promoter.

8. Figure 8 is a longitudinal section of a 30 day old transgenic rice grain showing transcriptional activity of the construct of Figure 7 (i.e. pLtp1.787-GN) containing the promoter of Figure 5. It is to be noted that transcriptional activity is achieved in the scutellar epithelial layer, as denoted by the blue staining.

5

9. Figure 9 is a longitudinal section of a mature germinating transgenic rice grain showing transcriptional activity of the construct of Figure 7 (i.e. pLtp1.787-GN) containing the promoter of Figure 5.

10 It is to be noted that transcriptional activity is achieved in the scutellar epithelial layer. Transcriptional activity is also observed in the shoot epithelial layer and in the aleurone layer. However, the extent of expression in the last two tissue types is not as pronounced as that in the scutellar epithelial layer.

15 However, more importantly, with the transgenic rice transcriptional activity is observed in the vascular tissue of the germinating seedling and the vascular tissue of the root and stem.

C. SUMMATION

20

The Examples relate to the isolation of and to the use of a 787 bp fragment of the promoter for the barley Ltp1 gene, which encodes a 10 kDa nsLTP. The gene was isolated by the use of a cDNA from a differential screening experiment in which the positive probe was constructed from aleurone cell poly (A) rich RNA, and the
25 negative probe from the starchy endosperm of immature grains.

A construct comprising the Ltp1 gene promoter fragment and a GOI (in this case *GUS*) was stably inserted into rice protoplasts.

30 Expression and *in situ* analysis for the wild type gene promoter demonstrated that the Ltp1 transcript is expressed in high levels only in the aleurone cells in developing barley caryopsis. This expression continued in germinating grains and also in

plantlets and mature plants.

However, for transgenic cereals, especially rice, even though there is some expression in the aleurone layer for the modified Ltp1 gene promoter it is, however, not as pronounced as that in each of the epithelial cells of the scutellum, the epidermal cells
5 of the coleoptile and the vascular strands of the embryo of developing caryopsis (or grain).

This result is completely unexpected as it shows that a modified Ltp1 promoter can
10 function differently in transgenic cereals, especially rice, than the wild-type Ltp1 gene in barley.

Expression and histochemical analysis for the transgenic rice demonstrated that the Ltp1 transcript is expressed in high levels in the scutellar epithelial tissue and vascular
15 tissue, especially of a germinating rice seedling and a developing rice grain and a rice plant (e.g. in the root, leaves and stem). This expression continued in germinating grains and also in plantlets and mature plants.

Importantly, for rice, expression is observed in the vascular tissue of the germinating
20 seedling and the vascular tissue of the root and stem.

This result is completely unexpected in view of the expression pattern of wild-type Ltp1 gene in barley.

25 Using the 787 bp promoter fragment in particle bombardments of developing barley caryopsis, we obtained activity (blue spots) in the epithelium layer of the scutellum.

The results therefore indicate that the modified Ltp1 gene promoter directs expression of a GOI predominantly in the aleurone cells of developing caryopsis, particularly for
30 barley, or the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant (e.g. in the root, leaves and stem) particularly for rice.

The modified Ltp1 gene promoter therefore represents a valuable tool for the expression of GOIs in the aleurone layer of developing caryopsis, in particular developing barley caryopsis.

- 5 Moreover, the modified Ltp1 gene promoter represents a valuable tool for the expression of GOIs in the scutellar epithelial cells and vascular cells of germinating seedlings or developing grain, in particular developing or germinating rice seedlings or grain. The epithelial or vascular expression is of particular benefit because the 787 bp LTP1 gene fragment can be used to express antisense α -amylase in the
10 scutellar epithelial layer in order to reduce or to prevent damage due to preharvest sprouting or to introduce or enhance pathogen resistance.

- One possible reason for the expression activity of the modified Ltp1 gene promoter of the present invention may be the absence of "silencer" elements in the modified
15 gene promoter which prevent expression of the wild type gene in, for example, the scutellar epithelial layer and vascular cells. Accordingly, the term "modified" (as defined above) could include removal of any silencer elements from the wild type Ltp1 gene promoter.

- 20 Studies with the modified Ltp1 gene promoter having deletions from and between the myb and myc sites when fused to GN showed that the relative activity of the deleted modified Ltp1 gene promoter was less (in some cases 70% less) than the modified Ltp1 gene promoter which contains the myb and myc sites. Therefore, it is believed that the presence of the myb and myc sites are important for even higher levels of
25 expression of the modified Ltp1 promoter in at least protoplasts of at least rice.

- Accordingly the present invention also covers a method of enhancing the *in vivo* expression of a GOI in at least the aleurone layer of a developing caryopsis or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a
30 developing grain or a plant preferably of an embryo of a developing monocotyledon grain or caryopsis, comprising stably inserting into the genome of those cells a DNA construct comprising a modified Ltp1 gene promoter and a GOI, wherein in the

formation of the construct the modified Ltp1 gene promoter is ligated to the GOI in such a manner that each of the myb site and the myc site in the modified Ltp1 gene promoter is maintained substantially intact.

- 5 The present invention also covers the use of a myb site and a myc site in a modified Ltp1 gene promoter to enhance *in vivo* expression of a GOI in at least the aleurone layer of a developing caryopsis or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant, preferably of an embryo of a developing monocotyledon caryopsis or grain, wherein the modified Ltp1
10 gene promoter and the GOI are integrated into the genome of the monocotyledon.

Each of these aspects is applicable to the combination expression system.

D. CONCLUSIONS VIS-A-VIS THE SPECIFIC EXAMPLES

- 15
1. The barley Ltp1 gene encodes a protein homologous to the 10 kDa wheat lipid transfer protein.
 2. The wild type Ltp1 gene promoter is expressed in developing barley aleurone
20 cells.
 3. The modified Ltp1 gene promoter is transiently expressed in developing barley scutellar epithelial cells after particle bombardment.
 - 25 4. The modified Ltp1 gene promoter directs expression of the *GUS*-reporter gene in the scutellar epithelial cells of developing transgenic rice grains. However, more pronounced expression is observed in the vascular tissue of germinating seedlings and the root and stem of the transgenic rice plant.
 - 30 5. The modified Ltp1 gene promoter contains sequence elements implicated in the transcriptional control of cereal endosperm specific genes.

6. The modified Ltp1 gene promoter contains myb and myc sequence elements that are implicated in the level of transcription in cereal endosperm.

Other modifications of the present invention will be apparent to those skilled in the
5 art without departing from the scope of the invention.

REFERENCES

- Bosnes, M., Weideman, F. and Olsen, O.-A. (1992) Endosperm differentiation in barley wild-type and sex mutants. *Plant J.* **2**, 661-674.
- 5 De Silva *et al.* 1992 WO 92/18634
- Dellaporta, S.L., Greenblatt, I., Kermicle, J.L., Hick, J.B. and Wessler, S. (1988) Molecular cloning of the *R-nj* allele by transposon tagging with *Ac*. In *Chromosome structure and function: Impact of new concepts*, 18th Stadler Genetics Symposium (Gustafson, J.P. and Appels, R., eds.), New York: Plenum Press, pp. 263-282.
- 10 Dieryck, W., Gautier, M.-F., Lullien, V. and Joudrier, P. (1992) Nucleotide sequence of a cDNA encoding a lipid transfer protein from wheat (*Triticum durum* Desf.). *Plant Mol. Biol.* **19**, 707-709.
- 15 Fincher, G.B. (1989) Molecular and cellular biology associated with endosperm mobilization in germinating cereal grains. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **40**, 305-346.
- 20 Fleming, A.J., Mandel, T., Hofmann, S., Sterk, P., de Vries, S.C., and Kuhlemeier, C. (1992) Expression pattern of a tobacco lipid transfer protein gene within the shoot apex. *Plant J.* **2**, 855-862.
- 25 Gordon-Kamm, W.J., Spencer, T.M., Mangano, M.L., Adams, T.R., Daines, R.J., Start, W.G., O'Brien, J.V., Chambers, S.A., Adams, W.R., Willetts, N.G., Rice, T.B., Mackey, C.J., Krueger, R.W., Kausch, A.P. and Lemaux, P.G. (1990) Transformation of maize cells and regeneration of fertile transgenic plants. *Plant Cell* **2**, 603-618.
- 30

- Jacobsen, J.V., Knox, R.B. and Pyliotis, N.A. (1971) The structure and composition of aleurone grains in the barley aleurone layer. *Planta* **101**, 189-209.
- Jakobsen, K., Klemsdal, S., Aalen, R., Bosnes, M., Alexander, D. and Olsen, O.-A. (1989) Barley aleurone cell development: molecular cloning of aleurone-specific cDNAs from immature grains. *Plant Mol. Biol.* **12**, 285-293.
- Jefferson, R.A. (1987) Assaying chimeric genes in plants: the *GUS* gene fusion system. *Plant Mol. Biol. Rep.* **5**, 387-405.
- Kader, J.-C., Julienne, M. and Vergnolle, C. (1984) Purification and characterisation of a spinach-leaf protein capable of transferring phospholipids from liposomes to mitochondria or chloroplasts. *Eur. J. Biochem.* **139**, 411-416.
- Karrer, E.J., Litts, J.C and Rodriguez, R.L. (1991) Differential expression of α -amylase genes in germinating rice and barley seeds. *Plant Mol. Biol.* **16**, 797-805.
- Klemsdal, S.S., Hughes, W., Lonneborg, A., Aalen, R. and Olsen, O.-A. (1991) Primary structure of a novel barley gene differentially expressed in immature aleurone layers. *Mol. Gen. Genet* **228** 9-16.
- Koltunow, A.M., Truettner, J., Cox, K.H., Wallroth, M. and Goldberg, R.B. (1990) Different temporal and spatial expression patterns occur during anther development. *Plant Cell* **2**, 1201-1224.
- Kosugi, S., Ohashi, Y., Nakajima, K. and Arai, Y. (1990) An improved assay for β -glucuronidase in transformed cells: methanol almost completely suppresses a putative endogenous β -glucuronidase activity. *Plant Sci.* **70**, 133-140.
- Kvaale, A. and Olsen, O.-A. (1986) Rates of cell division in developing barley endosperms. *Ann. Bot.* **57**, 829-833.

- Lea, R., Tommerup, H., Svendsen, I. and Mundy, J. (1991) Biochemical and molecular characterization of three barley seed proteins with antifungal properties. *J. Biol. Chem.* 266, 1564-73.
- 5 Linnestad, C., Lönneborg, A., Kalla, R. and Olsen, O.-A. (1991) The promoter of a lipid transfer protein gene expressed in barley aleurone cells contains similar Myb and Myc recognition sites as the maize *Bz-McC* allele. *Plant Physiol.* 97, 841-843.
- 10 McClintock, B. (1978) Development of the maize endosperm as revealed by clones. In *The clonal basis of development* (Subtelny, S. and Sussex, I.M., eds.), New York: Academic Press, pp.217-237.
- Monnet, F.-P. (1990) *Ph.D thesis*. Universite des Sciences et Techniques du
15 Languedoc, Montpellier, France, pp. 121.
- Mundy, J. and Rogers, J. (1986) Selective expression of a probable amylase/protease inhibitor in barley aleurone cells: comparison to the barley amylase/subtilisin inhibitor. *Planta* 169, 51-62.
- 20 Murashige, T., and Skoog, F. (1962) A revised medium for rapid growth and bioassays with tobacco tissue cultures. *Physiol. Plant.* 15, 473-497.
- Old & S.B. Primrose (1993) *Principles of Gene Manipulation - An Introduction to*
25 *Genetic Engineering 4th Edition*. Pub. Blackwell Scientific Publications. Pages 360-363.
- Olsen, O.-A., Jakobsen, K.S. and Schmelzer, E. (1990) Development of barley aleurone cells: temporal and spatial patterns of accumulation of cell specific mRNAs.
30 *Planta* 181 462-466

- Olsen, O.-A., Potter, R.H. and Kalla, R. (1992) Histo-differentiation and molecular biology of developing cereal endosperm. *Seed Sci. Res.* **2**, 117-131.
- Paz-Ares, J., Ghosal, D., Wienand, U., Peterson, P.A. and Saedler, H. (1987)
5 The regulatory *c1* locus of *Zea mays* encodes a protein with homology to myb proto-oncogene products and with structural similarities to transcriptional activators. *EMBO J.* **6**, 3553-3558.
- Schmelzer, E., Jahnen, W. and Hahlbrock, K. (1988) *In situ* localization of light-
10 induced chalcone synthetase mRNA, chalcone synthetase, and flavonoid end products in epidermal cells of parsley leaves. *Proc. Natl. Acad. Sci. U.S.A* **85**, 2989-2993.
- Shah *et al.* [1986] *Science* **233** 478-81
- 15 Shimamoto, K., Terada, R., Izawa, T. and Fujimoto, H. (1989) Fertile transgenic rice plants regenerated from transformed protoplasts. *Nature* **338**, 274-276.
- Skriver, K., Leah, R., Müller-Uri, F., Olsen, F.-L. and Mundy, J. (1992)
20 Structure and expression of the barley lipid transfer gene *Ltp1*. *Plant Molecular Biology* **18** 585-589.
- Slakeski, N. and Fincher, G.B. (1992) Developmental regulation of (1-3,1-4)- β -glucanase gene expression in barley. *Plant Physiol.* **99**, 1226-1231.
- 25 Smith, L.M., Handley, J., Li, Y., Martin, H., Donovan, L. and Bowles, D.J. (1992) Temporal and spatial regulation of a novel gene in barley embryos. *Plant Mol. Biol.* **20**, 255- 266.
- Somssich, I.E., Schmelzer, E., Kawalleck, P. and Hahlbrock, K. (1988) Gene
30 structure and *in situ* transcript localization of the pathogenesis-related protein 1 in parsley. *Mol. Gen. Genet.* **213**, 93-98.

Sossountzov, L., Riuz-Avila, L., Vignois, F., Jolliot, A., Arondel, V., Tchang, F., Grosbois, M., Guerbette, F., Miginiac, E., Delsney, M., Puigdomenech, P. and Kader, J.-C. (1991) Spatial and temporal expression pattern of a maize lipid transfer protein gene. *Plant Cell* 3, 923-933.

5

Sterk, P., Booij, H., Schellekens, G.A., Van Kammen, A. and De Vries, S.C. (1991) Cell-specific expression of the carrot EP2 lipid transfer protein gene. *Plant Cell* 3, 907-921.

- 10 Thoma, S., Kaneko, Y. and Sommerville, C. (1993) A non-specific lipid transfer protein from Arabidopsis is a cell wall protein. *The Plant Journal* 3(3), 427-436.

- Watanabe, S. and Yamada, M. (1986) Purification and characterization of a non-specific lipid transfer protein from germinated castor bean endosperms which transfers phospholipids and galactolipids. *Biochim. Biophys. Acta.* 876, 116-123.
- 15

SEQUENCE LISTING

(1) GENERAL INFORMATION

5 NAME OF APPLICANTS: O.-A. OLSEN AND R. KALLA
 BUSINESS ADDRESS: PLANT MOLECULAR BIOLOGY LABORATORY
 DEPARTMENT OF BIOTECHNICAL SCIENCES
 AGRICULTURAL UNIVERSITY OF NORWAY
 AND AGRICULTURAL BIOTECHNOLOGY PROGRAM NRC
 10 NORWAY
 N-1432

TITLE OF INVENTION: PROMOTER

15 (2) INFORMATION FOR SEQUENCE I.D. 1

SEQUENCE TYPE: NUCLEIC ACID
 MOLECULE TYPE: DNA (GENOMIC)
 ORIGINAL SOURCE: BARLEY
 20 SEQUENCE LENGTH: 787
 STRANDEDNESS: DOUBLE
 TOPOLOGY: LINEAR
 SEQUENCE:

25	-787	GAGCTCC	AAGGCATCAC	CAAGCTTCTA	TGACGCCAAA
	-750	ACATCCAAGA	AAGATATGTA	CTAGGATACC	AAGCACCCAA
		GAGTAAACGG	AGGAAGTATA	ATATAAGGCC	CTGTTTGATA
	-670	ACAAAGTAGT	AAAAAACTA	AAGTATTAAA	AACTGCAGTA
		ATTTTACGTG	TAGATAGAAA	ATACCATGGT	TTTAATATAA
30	-590	TAATATTTTT	TGCAGTATTC	ACAATGTAGA	GAAACTGTTT
		GATTACGCCA	CATATTACTG	CAGTTTAGAT	CGAGCAAGTA
	-510	CACGGGAAGA	AGATAACGAC	GTCCCACCCC	TTCTTTTCGC
		CTTCTCTGTT	TTTTAAAAAG	AGGTCTGGGG	TTAGTTTTTT
	-430	CAATACTGCA	GTTTTAAAAT	CACAATTCTT	AGAGGCAACC
35		AAACACCTCA	TTGTAAATAA	AACTATGATA	ATCTCCAAAA
	-350	CTGCAGTATT	CTAAAAATAC	TACAAAAATT	CTTTGTTATC
		AAACAGGGCC	TAAGGAGTTA	AAAAAATTTA	GCCGTAACGT
	-270	AGACTCGGCG	AGGCACCAGC	AGCTAGCAGT	CATCAACACT
		TGATGGTTGG	CAAAGCCGAG	TCGACGTGTC	GCGGGGCTCG
40	-190	GCCTGAGCGG	GAGATACAAT	CTGTTCTCCA	GTAACCCCGT
		CGATTTGGCC	CGCCGACTAA	AGCATCCAGG	CATCTCTCGC
	-110	TCGAACCCCT	ATTTAAGCCC	CTCCATTCTT	CCCAACATTC
		TCCACACCTC	CACGAGTTGC	TCATCACTAG	CTAGTACGTT
45	-30	GTA CTGTTAG	CTACAGATTA	AGAAGTGATC	

NOTE: ABOVE SEQUENCE IS A RETYPED VERSION OF FIGURE 5 WHICH IS TO BE
 TAKEN AS THE CORRECT SEQUENCE

(3) INFORMATION FOR SEQUENCE I.D. 2

	SEQUENCE TYPE:	NUCLEIC ACID
	MOLECULE TYPE:	DNA (GENOMIC)
5	ORIGINAL SOURCE:	BARLEY
	SEQUENCE LENGTH:	-807
	STRANDEDNESS:	DOUBLE
	TOPOLOGY:	LINEAR
	SEQUENCE:	

10	-807	GATCTCG	ATGTGTAGTC	TACGAGAAGG
	-780	TCTTCGTGAG	AATAACCGTG	GCCTAAAAAT
		GGATAAATAA	AATGTGGTGG	TACAGTACTT
		ACTCATCAAG	AGGATGCTTT	TCCGATGAGC
15	-660	CATCGGACCT	CACATACCTC	CATTGTGGTG
		TGCTCATTTA	GTGATGGGTA	AATTTTGTIT
		AGGTTTTGAC	ATTTCAGTTT	TGCCACTCTT
	-540	AAATAATTTT	CATTCCGCGG	CAAAGCAAA
		TTTACTTTTA	CCACTCTTAG	CTTTCACAAT
20		TGCCACTCTA	GAAATTCTGT	TTATGCCACA
	-420	AAAAACACTC	ACTTATTTGA	AGCCAAGGTG
		GGAAATGTGA	CATAAAGTAA	CGTTCGTGTA
		TTGTACTCCT	CGTAACAAGA	GACGGAACA
	-300	ATCGCGTTTG	GAAGGCTTTG	CATCACCTTT
25		CATGAATGGA	GTCGTCTGCT	TGCTAGCCTT
		CCACTGAGTC	CGGGCGGCAA	CTACCATCGG
	-180	AGCTGACCTC	TACCGACCGG	ACTTGAATGC
		TCAGCGACGA	TGGCCGCGTA	CGCTGGCGAC
		ATGCATGGCG	GCACATGGCG	AGCTCAGACC
30	-60	GGCTACAAAT	ACGTACCCCG	TGAGTGCCCT
		TTACACCTGC		

43

INDICATIONS RELATING TO A DEPOSITED MICROORGANISM

(PCT Rule 13bis)

A. The indications made below relate to the microorganism referred to in the description on page <u>24</u> , lines <u>15-17</u>	
B. IDENTIFICATION OF DEPOSIT Further deposits are identified on an additional sheet <input type="checkbox"/>	
Name of depositary institution The National Collections of Industrial and Marine Bacteria Limited (NCIMB)	
Address of depositary institution (including postal code and country) 23 St. Machar Drive Aberdeen Scotland AB2 1RY United Kingdom	
Date of deposit <div style="text-align: right;">11. JAN. 1994</div>	Accession Number <div style="text-align: right;">NCIMB 40609</div>
C. ADDITIONAL INDICATIONS (leave blank if not applicable) This information is continued on an additional sheet <input type="checkbox"/>	
<p>In respect of those designations in which a European patent is sought, and any other designated state having equivalent legislation, a sample of the deposited microorganism will be made available until the publication of the mention of the grant of the European patent or until the date on which the application has been refused or withdrawn or is deemed to be withdrawn, only by the issue of such a sample to an expert nominated by the person requesting the sample. (Rule 28(4) EPC).</p>	
D. DESIGNATED STATES FOR WHICH INDICATIONS ARE MADE (if the indications are not for all designated States)	
E. SEPARATE FURNISHING OF INDICATIONS (leave blank if not applicable)	
The indications listed below will be submitted to the International Bureau later (specify the general nature of the indications e.g., "Accession Number of Deposit")	

<div style="border-bottom: 1px solid black; padding-bottom: 5px;"> For receiving Office use only </div> <div style="border: 1px solid black; padding: 5px;"> <input type="checkbox"/> This sheet was received with the international application </div> <div style="border: 1px solid black; padding: 5px; margin-top: 10px;"> Authorized officer </div>	<div style="border-bottom: 1px solid black; padding-bottom: 5px;"> For International Bureau use only </div> <div style="border: 1px solid black; padding: 5px;"> <input type="checkbox"/> This sheet was received by the International Bureau on: </div> <div style="border: 1px solid black; padding: 5px; margin-top: 10px;"> Authorized officer </div>
--	---

CLAIMS

1. A modified Ltp1 gene promoter which is integrated, preferably stably integrated, within a plant material's genomic DNA and which is capable of inducing
5 expression of a GOI when fused to the gene promoter in at least the aleurone cells or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant (e.g. in the root, leaves and stem).
2. A modified Ltp1 gene promoter according to claim 1 wherein the plant
10 material is a developing caryopsis, a germinating seedling, a developing grain or a plant and wherein the gene promoter is integrated, preferably stably integrated, in the developing caryopsis's genomic DNA or the germinating seedling's genomic DNA or the developing grain's genomic DNA or the plant's genomic DNA and which is capable of inducing expression of a GOI when fused to the gene promoter in at least
15 the aleurone cells of the developing caryopsis or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant (e.g. in the root, leaves and stem).
3. A modified Ltp1 gene promoter according to claim 1 or claim 2 wherein the
20 promoter comprises the nucleic acid sequence shown as SEQ. I.D. 1, or a sequence that has substantial homology with that of SEQ. I.D. 1, or a variant thereof.
4. An isolated Ltp1 gene promoter comprising the sequence shown as SEQ. I.D.
1, or a sequence that has substantial homology therewith, or a variant thereof.
25
5. A construct comprising

a GOI and

30 a modified Ltp1 gene promoter according to any one of claims 1 to 4;

wherein the construct is capable of being expressed in at least the aleurone cells or in at least the scutellar epithelial tissue or vascular tissue of a plant material; and

5 wherein if there is expression in just the aleurone layer of a developing barley caryopsis then the fused promoter and GOI are not the 769 bp fragment of Skriver *et al* (1992).

6. A construct according to claim 5 wherein the construct is capable of being
10 expressed in at least the aleurone cells of a developing caryopsis or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant when the construct is integrated, preferably stably integrated, within the caryopsis's or grain's or seedling's or plant's genomic DNA.

15 7. A construct according to claim 5 or claim 6 wherein the modified Ltp1 gene promoter comprises the nucleic acid sequence shown as SEQ. I.D. 1, or a sequence that has substantial homology with that of SEQ. I.D. 1, or a variant thereof.

8. The construct according to any one of claims 5 to 7 wherein the construct
20 further comprises at least one additional sequence to increase expression of the GOI.

9. An expression system for at least the aleurone cells or for at least the scutellar epithelial tissue or vascular tissue of a plant material, the expression system comprising

25

a GOI fused to a modified Ltp1 gene promoter

wherein the expression system is capable of being expressed in at least the aleurone cells or in at least the scutellar epithelial tissue or vascular tissue of the plant
30 material; and

wherein if there is expression in just the aleurone layer of a developing barley caryopsis then the fused promoter and GOI are not the 769 bp fragment of Skriver *et al* (1992).

- 5 10. An expression system according to claim 9 wherein the expression system is for at least the aleurone cells of a developing caryopsis or for at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or developing grain or plant (e.g. in the root, leaves and stem).
- 10 11. An expression system according to claims 9 or claim 10 wherein the expression system is additionally capable of being expressed in the embryo cells of the germinating grain or the plantlet.
12. An expression system according to any one of claims 9 to 11 wherein the
15 expression system is integrated, preferably stably integrated, within a developing caryopsis's genomic DNA or a germinating seedling's genomic DNA or a developing grain's genomic DNA or a plant's genomic DNA.
13. An expression system according to any one of claims 9 to 12 wherein the gene
20 promoter comprises the sequence shown as SEQ I.D. No. 1 or comprises a sequence that has substantial homology therewith, or is a variant thereof.
14. An expression system for at least the aleurone cells of a developing caryopsis or for at least the scutellar epithelial tissue or vascular tissue of a germinating
25 seedling or developing grain or plant (e.g. in the root, leaves and stem), the expression system comprising

a gene promoter fused to a GOI

- 30 wherein the expression system is capable of being expressed in at least the aleurone cells of the developing caryopsis or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant (e.g. in

the root, leaves and stem); either

wherein if there is expression in just the aleurone layer of a developing barley caryopsis then either the promoter is not the wild type Ltp1 promoter in its natural
5 environment and the GOI is not the Ltp1 functional gene in its natural environment; or

wherein if there is expression in just the aleurone layer of a developing caryopsis then the fused promoter and GOI are not the 769 bp fragment of Skriver
10 *et al* (1992).

15. An expression system according to any one of claims 9 to 14 comprising a construct according to any one of claims 5 to 8.

16. A transgenic cereal comprising an expression system according to any one of
15 claims 9 to 15 or a construct according to any one of claims 5 to 8 wherein the expression system or construct is integrated, preferably stably integrated, within the cereal's genomic DNA.

17. The use of a gene promoter as defined in any one of the preceding claims to
20 induce expression of a GOI when fused to the gene promoter in at least the aleurone cells or in at least the scutellar epithelial tissue or vascular tissue of a plant material.

18. The use according to claim 17 wherein the gene promoter is used to induce
25 expression of a GOI when fused to the gene promoter in at least the aleurone cells of a developing caryopsis or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant (e.g. in the root, leaves and stem).

19. A process of expressing a GOI when fused to a gene promoter as defined in
30 any one of the preceding claims, wherein expression occurs in at least the aleurone cells or in at least the scutellar epithelial tissue or vascular tissue of a plant material.

20. A process according to claim 19 wherein the gene promoter expresses the GOI when fused to the gene promoter in at least the aleurone cells of a developing caryopsis or in at least the scutellar epithelial tissue or vascular tissue of a germinating seedling or a developing grain or a plant (e.g. in the root, leaves and stem).

21. A process according to claim 19 or claim 20 wherein the promoter and GOI are integrated, preferably stably integrated, within a cereal's genomic DNA.

22. A process of expressing in at least the scutellar epithelial tissue or vascular tissue of a developing grain or a germinating seedling or a plant, preferably a developing rice grain or a germinating rice seedling or a transgenic rice plant, an expression system according to any one of claims 9 to 15 or a construct according to any one of claims 5 to 8 wherein the expression system or construct is integrated, preferably stably integrated, within the cereal's genomic DNA.

23. The invention of any one of claims 1 to 22 wherein the gene promoter is a fragment of a barley Ltp1 gene promoter.

24. The invention of claim 23 wherein the promoter is for a 10 kDa lipid transfer protein.

25. The invention of claim 23 or claim 24 wherein the gene promoter is obtainable from plasmid NCIMB 40609.

26. The invention of any one of claims 1 to 15 wherein the gene promoter is used for expression of a GOI in a cereal caryopsis or a cereal grain or a cereal seedling or a cereal plant.

27. The invention of claim 26 wherein the cereal caryopsis is a developing cereal caryopsis, the cereal grain is a developing cereal grain, and the cereal seedling is a germinating cereal seedling.

28. The invention of claim 26 or claim 27 wherein the cereal is any one of a rice, maize, wheat, or barley.

29. The invention of claim 28 wherein the cereal is rice or maize, preferably rice.

5

30. The invention according to any one of claims 1 to 29 wherein the developing caryopsis is a developing barley caryopsis, the germinating seedling is a germinating rice seedling, the developing grain is a developing rice grain, and the plant is a transgenic rice plant.

10

31. A combination expression system comprising

a. as a first construct, a construct according to any one of claims 5 to 8; and

15 b. as a second construct, a construct comprising a GOI and another gene promoter that is tissue- or stage-specific.

32. A combination expression system according to claim 31 wherein each construct is integrated, preferably stably integrated, within a plant material.

20

33. A combination expression system according to claim 32 wherein each construct is integrated, preferably stably integrated, within a developing caryopsis's genomic DNA or a grain's genomic DNA or a seedling's genomic DNA or a plant's genomic DNA.

25

34. A combination expression system according to any one of claims 31 to 33 wherein the first construct comprises a modified Ltp1 gene promoter comprising the nucleic acid sequence shown as SEQ. I.D. 1, or a sequence that has substantial homology with that of SEQ. I.D. 1, or a variant thereof.

30

35. A combination expression system according to any one of claims 31 to 34 wherein the promoter in the second construct is an aleurone specific promoter.

36. A combination expression system according to any one of claims 31 to 35 wherein the promoter in the second construct is a barley promoter.
37. A combination expression system according to any one of claims 31 to 35
5 wherein the second construct is the B22E gene promoter.
38. A combination expression system according to any one of claims 31 to 37 wherein the promoter in the second construct is the Ltp2 gene promoter.
- 10 39. A combination expression system according to claim 38 wherein the promoter in the second construct is for a 7 kDa lipid transfer protein.
40. A combination expression system according to claim 38 or 39 wherein the promoter in the second construct is the promoter for Ltp2 of *Hordeum vulgare*.
15
41. A combination expression system according to any one of claims 31 to 40 wherein the promoter in the second construct comprises the sequence shown as SEQ. I.D. 2, or a sequence that has substantial homology therewith, or a variant thereof.
- 20 42. A combination expression system according to any one of claims 38 to 41 wherein each of the myb site and the myc site in the Ltp2 gene promoter is maintained substantially intact.
43. A combination expression system according to any one of claims 31 to 42
25 wherein the second construct further comprises at least one additional sequence to increase expression of the GOI.
44. A developing cereal grain, preferably a germinating rice seedling, comprising any one of: a promoter according to any one of claims 1 to 4 or any claim dependent thereon, an expression system according to any one of claims 9 to 15 or any claim dependent thereon, a construct according to any one of claims 5 to 8 or any claim dependent thereon, or a combination expression system according to any one of
- 30

claims 31 to 43 or any claim dependent thereon.

45. The invention of any one of the preceding claims wherein each of the myb site and the myc site in the gene promoter is maintained substantially intact.


5

46. Plasmid NCIMB 40609.

47. A promoter, a construct or an expression system or a combination expression system substantially as described herein with reference to any one of Figures 5 to 9.

10

Fig 1



2 / 1 0

FIG 2




Fig 3

1 GAGCTCCAAGGCATCACCAAGCTTCTATGACGCCAAAACATCCAAGAAAGATATGTACTAGGATACCAAGCACCC
76 AAGAGTAAACGGAGGAAGTATAATATAAGGCCCTGTTTGATAACAAAGTAGTAAAAAACTAAAGTATTAACAAAC
151 TGCAGTAAATTTACGTGTAGATAGAAAATACCATGGTTTTAATATAATAATATTTTTTGCAGTATTCACAATGTA
226 GAGAACTGTTTGATTACGCCACATATTACTGCAGTTTAGATCGAGCAAGTACACGGGAAGAAGATAACGACGTC
301 CCACCCCTTCTTTTCGCCTTCTCTGTTTTTAAAAAGAGGTCTGGGGTTAGTTTTTCAATACTGCAGTTTTAAA
376 ATCACAATTCTTAGAGGCAACCAACACCTCATTGTAATAAACTATGATAATCTCCAAAACGTCAGTATTCTA
451 AAAATACTACAAAAATTCTTTGTTATCAACAGGGCCTAAGGAGTTAAAAAAATTTAGCCGTAACCTGAGACTCGG
526 CGAGGCACCCAGCAGCTAGCAGTCATCAACACTTGATGGTTGGCAAAGCAGAGTCGAGCTGTCGCGGGGCTCGGCC
601 TGAGCGGGAGATCAATCTGTTCTCCAGTAACCCCGTCGATTGGCCCCGCCGACTAAAGCATCCAGGCATCTCTC
676 GCTCGAACCCCTATTAAACCCCTCCATTCTCCCAACATTCTCCACACCTCCACGAGTTGCTCATCACTAGCTA
751 GTACGTTGTACTGTTAGCTACAGATTAAGAAGTGATC ATG GCC CGC GCT CAG GTA CTG CTC ATG
M A R A Q V L L M
815 GCC GCC GCC TTG GTG CTG ATG CTC ACG GCG GCC CCG CGC GCT GCC GTG GCC CTC AAC
A A A L V L M L T A A P R A A V A L N
872 TGC GGC CAG GTT GAC AGC AAG ATG AAA CCT TGC CTG ACC TAC GTT CAG GGC GGC CCC
C G Q V D S K M K P C L T Y V Q G G P
929 GGC CCG TCC GGC GAA TGC TGC AAC GGC GTC AGG GAT CTC CAT AAC CAG GCG CAA TCC
G P S G E C C N G V R D L H N Q A Q S
986 TCG GGC GAC CGC CAA ACC GTT TGC AAC TGC CTG AAG GGC ATC GCT CGC GGC ATC CAC
S G D R Q T V C N C L K G I A R G I H
1043 AAT CTC AAC CTC AAC AAC GCC GCC AGC ATC CCC TCC AAG TGC AAT GTC AAC GTC CCA
N L N L N N A A S I P S K C N V N V P
1100 TAC ACC ATC AGC CCC GAC ATC GAC TGC TCC AG gtgattaaatttacactcatccagagtgaat
Y T I S P D I D C S R
1164 ctttaaaaagaactatatatttacgaacggagtgagtatataggaacattcatccacgtaaaatttggtgatattaa
1240 cattaacacgcatgattgacctgcag G ATT TAC TGAGCGACGATCCGTCAAGCTGGTGCTCAGCTCATCGA
I Y *
1310 TCCACGTGGAGCTGAAGCGCGCAGCCTCTGTCCCTATGTAGTATGGCTACCAGTTATGCCGAGTTTATGCTGAAT
1385 AAGAACTCTCTCCTGTACTCCTTTGGAGGAGATCAGTATCTATGTACGTGAGAGTTGAGAGTTTGTACCATCGGC
1460 ACTCCCAGTGTATTATGGACTATATGCAT

Fig 4

GTCCACAACCTCATGAGCATCACGGAATGGCATGAGTTGAAATATAACTACATTGCTCAAA -1621
GCAACAAAAAGCACATTAGAATCTTGAGCATTGAGATAAGAGTTTTTCTCATGCTCTAAA -1561
TATATATTTTGAGAATCCTTTGGAGGAGAAAAATCCATATTTACAATTCGTTGTAAATTT -1501
GAGTCCATGATCCTAAAGAGATTAAGCATGCGAATTACCCAAACATCAAAATTTGTGCCA -1441
TTGAAACTAAGAGTGTTAGAGAATCCTAATCCCCTAGTTGACATACTTACTCTCTAGGTG -1381
GTGAAACCTAATAATGAGAGATCTAGCTCTAATACCAATTGAGAGGATGTGGATGTCGCC -1321
TAGAGGGGGCGGTGAATAGGCGCTTTAAAATAATTACGGTTTAGGCTCGAACAAATGTGGA -1261
ATAAACTAACGTTTCATTTGTCAAGCGCAAAACCTAAAACAACCTAGGCTCACCTATGTG -1201
CACCAACAACCTTATGATAAGCAAGATAAAAAAACTAAGTGATGGCAGAATATATAACAAG -1141
AAACAATATGGCTATCACAAAGTGAAGTGCATAAGTAAACAGCTCGGGTAAGGGACAACC -1081
GAGCCATGCGGAGACGACGATGTATCCTCAAGTTCACACACTTGCGGATGCTAATCTCCG -1021
TTTGAAGCAGTGTGGAGGCACAATCGTCCCCAAGAAGCCACTAAGGCCACCGTAATCTCC -961
TCACGCCCTCGCACAATCGAAGATGTTGTGATTCCACTAAGGGACCCCTTGAGGGCAGTCA -901
CTGAACCCGTATAAACATGGTTGGAACAATCTCCACGACTTAATTGGAGACTCCCAACAA -841
CACCACGAACCTTCATCATAACGAAATATGGCTTCGAGGTAACCTCAAATGCTCGGGGCA -781
ATTTTACAACCTAATTGAAGACCTCGACGCTTGCGTGGAGCTTTACACTATAATGATTG -721
AGCTCCAAGGGCATCACCAAGCTTCTATGACGCCAAAACATCCAAGAAAGATATGTACTA -661
GGATACCAAGCACCCCAAGAGTAAACGGAGGAAGTATAATATAAGGCCCTGTTTGATAACA -601
AAGTAGTAAAAAACTAAAGTATTAAAACTGCAGTAATTTTACGTGTAGATAGAAAATA -541
CCATGGTTTAAATATAATAATATTTTTTGCAGTATTCAATGTAGAGAACTGTTTGAT -481
TACGCCACATATTACTGCAGTTTAGATCGAGCAAGTACACGGGAAGAAGATAACGACGTC -421
CCACCCCTTCTTTTCGCCTTCTCTGTTTTTTAAAAAGAGGTCTGGGGTTAGTTTTTTCAA -361
TACTGCAGTTTTAAATCACAATTCCTAGAGGCAACCAACACCTCATTGTAAATAAAAC -301
TATGATAATCTCCAAAACCTGCAGTATTCTAAAAATACTACAAAATTCCTTTGTATCAAA -241
CAGGGCCTAAGGAGTTAAAAAAATTTAGCCGTAACCTGAGACTCGGCGAGGCACCAGCAGC -181

Fig 4 contd.


TAGCAGTCATCAACACTTGATGGTTGGCAAAGGCGAGTCGACGTGTCGCGGGGCTCGGCC -121
 TGAGCGGGAGATACAATCTGTTCTCCAGTAACCCCGTCGATTGGCCCCGCCGACTAAAGC -61
 ATCCAGGCATCTCTCGCTCGAACCCCTATTAAAGCCCCCTCCATTCTCTCCCAACATTCTCC -1
 ACACCTCCACGAGTTGCTCATCACTAGCTAGTACGTTGTACTGTTAGCTACAGATTAAGA 60
 AGTGATCATGGCCCCGCGCTCAGGTACTGCTCATGGCCGCGCCTTGGTGCTGATGCTCAC 120
 M A R A Q V L L M A A A L V L M L T
 GCGGCCCCCGCGCTGCGGTGGCCCTCAACTGCGGCCAGGTTGACAGCAAGATGAAACC 180
 A A P R A A V A L N C G Q V D S K M K P
 TTGCCTGACCTACGTTTCAGGGCGGCCCCGCGCCGTCGCGGGAATGCTGCAACGGCGTCAG 240
 C L T Y V Q G G P G P S G E C C N G V R
 GGATCTCCATAACCAGGCGCAATCCTCGGGCGACCGCCAAACGTTTGCAACTGCCTGAA 300
 D L H N Q A Q S S G D R Q T V C N C L K
 GGGGATCGCTCGCGGCATCCACAATCTCAACCTCAACAACGCCGCCAGCATCCCTCCAA 360
 G I A R G I H N L N L N N A A S I P S K
 GTGCAATGTCAACGTCCCATACACCATCAGCCCCGACATCGACTGCTCCAGgtgattaaa 420
 C N V N V P Y T I S P D I D C S R
 ttacactcatccagagtgaatcttttaaaagaactatatatttacgaacggagtgagtat 480
 ataggaacattcatccacgtaaaaatttggtgatattaacattaacacgcatgattgacct 540
 gcagGATTTACTGAGCGACGATCCGTCAGCTGGTGCTCAGCTCATCGATCCACGTGGAG 600
 I Y *
 CTGAAGCGCGCAGCCTCTGTCCCTATGTAGTATGGCTACCAGTTATGCCGAGTTTATGCT 660
 GAATAAGAACTCTCTCCTGTACTCCTTTGGAGGAGATCAGTATCTATGTACGTGAGAGTT 720
 GAGAGTTTGTACCATCGGCACTCCAGTGTTTATGGACTATATGCATACACCTCCTTCTG 780
 TGCTCAGTGTGTAACCTTGTCTCTCTGTTTCCTCACGTTCCGCTCTCATATAATAATTTAC 840
 TTATGTGCTCTAGGATCGTAGTACAGTATCATATATATACCTCTCTATGAATTAGTTTAC 900
 CGTAGACCGTATGTTTCTTGAATCTGGATGAAAATTACGGATTCAAGCGTGCGTCCCGCA 960
 TATAATAAGCTTGCTTACGGATTCAAGCGTGCGTCACGCGGCTCAGTAGATGATGAGGAT 1020
 ACTCGCTGCTGCATCTCTACATCCCGCTCATGAGCTGAGCTGAGCCCGGGTCTCCCCCG 1080
 CTCCGCCCCGCTGGCCACCCCGCGCGCGCCGACCCTCAAACAGCCTTCATGACGAGCCGCC 1140
 CGCCAGCAAGATCTGTTGGCTCCTCCCTGTCCGTCGTAGAGAAACCCAGCA 1192

Fig 5

1 GAGCTCCAAGGCATCACCAAGCTTCTATGACGCCAAAACATCCAAGAAAGATATGTACTAGGATACCAAGCACCC
76 AAGAGTAAACGGAGGAAGTATAATATAAGGCCCTGTTTGATAACAAAGTAGTAAAAAACTAAGTATTAAAAAC
151 TGCAGTAATTTACGTGTAGATAGAAAATACCATGGTTTTAATATAATAATATTTTTTGCAGTATTCACAATGTA
226 GAGAAACTGTTTGATTACGCCACATATTACTGCAGTTTAGATCGAGCAAGTACACGGGAAGAAGATAACGACGTC
301 CCACCCCTTCTTTTCGCCCTTCTCTGTTTTTAAAAAGAGGTCTGGGGTTAGTTTTTCAATACTGCAGTTTTTAAA
376 ATCACAATTCCTAGAGGCAACCAACACCTCATTGTAAATAAACTATGATAATCTCCAAAAGTGCAGTATTCTA
451 AAAATACTACAAAAATTCTTTGTTATCAAACAGGGCCTAAGGAGTTAAAAAAATTTAGCCGTAAGTGCAGTCTGG
526 CGAGGCACCAGCAGCTAGCAGTCATCAACACTTGATGGTTGGCAAAGCAGACTCGACGTGTCGCGGGGCTCGGCC
601 TGAGCGGGAGATACAACTGTTCTCCAGTAACCCCGTCGATTTGGCCCGCCGACTAAAGCATCCAGGCATCTCTC
676 GCTCGAACCCCTATTTTAAACCCCTCCATTCTCCCAACATTCTCCACACCTCCACGAGTTGCTCATCACTAGCTA
751 GTACGTTGTACTGTTAGCTACAGATTAAGAAGTGATC

7 / 1 0

Fig 6



8 / 1 0

Fig. 7




Fig. 8





Fig. 9



INTERNATIONAL SEARCH REPORT

Inter. Application No
PC./NO 95/00042

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C12N15/82 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
O,X	J. CELL. BIOCHEM. SUPPL. 0, 1994 page 99 O.-A. OLSEN ET AL.; 'The barley LTP2 promoter yields high level of GUS expression in the aleurone layer of developing grains of transgenic rice' see abstract no. X1-213. & Keystone Symposium on improved crop and plant products through biotechnology, Keystone, Colorado, USA, January 9-16, 1994. --- -/--	1-21, 25-30, 44-47

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

* Special categories of cited documents :

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

30 June 1995

Date of mailing of the international search report

17.07.95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Yeats, S

INTERNATIONAL SEARCH REPORT

International Application No

PC1/NO 95/00042

C(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>PLANT MOL. BIOL., vol. 18, no. 19, 1992 page 585-589 K. SKRIVER ET AL.; 'Structure and expression of the barley lipid transfer protein gene Ltpl' cited in the application see the whole document. ---</p>	<p>1-21, 23-30, 44,45,47</p>
X	<p>PLANT J., vol. 2, 1992 pages 855-862, A.J. FLEMING ET AL.; 'Expression pattern of a tobacco lipid transfer protein gene within the shoot apex' cited in the application see the results section. ---</p>	<p>1-22,47</p>
X	<p>PLANT CELL, vol. 3, 1991 pages 923-933, L. SOSSOUNTZOV ET AL.; 'Spatial and temporal expression of a maize lipid transfer protein gene' see pages 923-925. ---</p>	<p>1-22, 26-29, 44,47</p>
A	<p>PLANTA, vol. 192, 1994 pages 574-580, K. GAUSING; 'Lipid transfer protein genes specifically expressed in barley leaves and coleoptiles' see the abstract, Figure 2 and page 577. ---</p>	<p>1</p>
X	<p>PLANT PHYSIOL., vol. 97, 1991 pages 841-843, C. LINNESTAD ET AL.; 'Promoter of a lipid transfer protein gene expressed in barley aleurone cells contains similar myb and myc recognition sites as the maize Bz-MyC allele' cited in the application see the whole document. -----</p>	<p>21,23, 24,26, 28,44, 45,47</p>